

**NOTES ON A CONFORMAL CHARACTERIZATION OF 2-DIMENSIONAL
LORENTZIAN MANIFOLDS WITH CONSTANT RICCI SCALAR
CURVATURE**

Nicolò Cangiotti¹ and Mattia Sensi²

We present a characterization of 2-dimensional Lorentzian manifolds with constant Ricci scalar curvature. It is well known that every 2-dimensional Lorentzian manifolds is conformally flat, so we rewrite the Ricci scalar curvature in terms of the conformal factor and we study the solutions of the corresponding differential equations. Several remarkable examples are provided.

Keywords: Lorentzian manifolds, 2-dimensional manifolds, Minkowski space, conformal factor, Ricci scalar curvature.

MSC2020: 35L05, 53B30, 83C15.

1. Introduction

Pseudo-Riemannian manifolds have historically been of interdisciplinary interest, and they have been extensively studied in many fields, from differential geometry to mathematical physics. In particular, there is a class which, for many reasons, has assumed an importance comparable to the one of Riemannian manifolds. It is indeed well known that, in the general theory of relativity, the spacetime has historically been modeled as a 4-dimensional Lorentzian manifold. Lorentzian manifolds represent a very interesting case study, which has been thoroughly investigated in mathematics [1, 2, 5] as much as in physics [9, 14, 18]. And, as it often happens in geometry, there are many properties which depend completely on the dimension of the manifold.

This paper fits into the body of works that propose an analysis of the 2-dimensional Lorentzian geometry (see, for instance, [4, 6, 8, 13, 17, 19]). Our idea is the following: starting from the fact that every 2-dimensional Lorentzian manifold is conformally flat, we show a very simple way to construct Lorentzian manifolds with constant Ricci scalar curvature by deriving different expressions of the conformal factor. In particular, we provide explicit forms for the conformal factor, which is a solution of a classical partial differential equations. In the same way, we propose also several examples, some that recover very familiar Lorentzian manifolds, whereas other are provided to illustrate the potential of our method.

The paper is organized as follows. In Section 2, we recall some definitions and a classical result regarding 2-dimensional Lorentzian manifolds. Section 3 is devoted to illustrate a way to construct Lorentzian manifolds starting from the conformal factor. Finally, we conclude with Section 4 by summarizing the ideas exhibited and by presenting possible further developments.

¹University of Pavia, Department of Mathematics, via Ferrata 5,, 27100 Pavia (PV), Italy. Email: nicolo.cangiotti@unipv.it

²University of Trento, Department of Mathematics, via Sommarive 14,, 38123 Trento (TN), Italy. Email: mattia.sensi@unitn.it

2. Definitions and basic concepts

In this section, we recall some basic definitions and concepts regarding Lorentzian manifolds; we remark similarities with the Riemannian case. Moreover, we state a fundamental theorem on the conformal structure of the 2-dimensional Lorentzian manifolds.

Definition 2.1. *A pseudo-Riemannian manifold \mathcal{M} is a differentiable manifold, equipped with an everywhere non-degenerate metric g . We denote a manifold provided with a specific metric with the couple (\mathcal{M}, g) .*

It is well known that, by Sylvester's law of inertia, one can identify the metric with its signature [12]. We restrict our study to a particular signature, which, as mentioned above, is very relevant in many physical context, especially in the field of the general theory of relativity.

Definition 2.2. *A Lorentzian manifold is a pseudo-Riemannian manifold \mathcal{M} , equipped with a metric g with signature $(1, n - 1)$, where n is the dimension of \mathcal{M} .*

For our purpose, we shall consider the 2-dimensional Lorentzian manifolds (it is appropriate to underline that the definitions below can easily be extended to the manifolds higher dimension). As does the Euclidean space in the case of Riemannian manifolds, a key role is played by the *Minkowski* space.

Definition 2.3. *Let us consider the following metric tensor η :*

$$\eta = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}.$$

A Lorentzian manifold \mathcal{M} , equipped with the metric η , is the Minkowski space. The metric η is called Minkowski metric.

Remark 2.1. *Historically, the definition of Minkowski space in the literature refers to the 4-dimensional space [9], but this little abuse of notation should not cause confusion, as we will work in the 2-dimensional case, unless explicitly stated otherwise.*

The next definition introduce the notion of *flat manifold*, by using the very usual concept of *Ricci scalar curvature*.

Definition 2.4. *A manifold (\mathcal{M}, g) with a null Ricci scalar curvature is called flat manifold. In this case, we shall call its metric flat metric. A manifold (\mathcal{M}, \tilde{g}) is conformally flat if its metric can be expressed, via the composition with a smooth function Ω , as*

$$\tilde{g} = \Omega \cdot g,$$

where g is a flat metric.

Remark 2.2. *It is clear that the Minkowski space of Def. 2.3 is a flat manifold.*

In the previous definitions, we implicitly express the metric in terms of a tensor (actually a matrix, in this case) as in Def. 2.3. However, it is possible to give a different, and very useful, characterization for the metric, namely the *line element*.

Definition 2.5. *Let us consider a metric g_{ij} , where we indicate rows and columns of the matrix associated to g by the indices i and j , respectively. By considering the general curvilinear coordinates, it is possible to define the line element ds^2 as:*

$$ds^2 = g_{ij} dq^i dq^j, \quad i, j = 1, 2.$$

Here we are using the Einstein summation convention.

Remark 2.3. For the metric η of Def. 2.3 the line metric for the coordinates (t, x) could be expressed by

$$ds^2 = -dt^2 + dx^2$$

The fundamental result which inspired this work is given by the following theorem, which gives a very powerful strategy to handle 2-dimensional Lorentzian manifolds.

Theorem 2.1. Every 2-dimensional Lorentzian manifold (\mathcal{M}, g) is conformally flat.

For the proof see, e.g., Theorem 7.2 in [16]. The next section is devoted to exploring the conformal factor in the case of manifolds with constant Ricci scalar curvature.

Remark 2.4. Kühnel and Rademacher produced a very interesting study regarding conformal geometry. In [10], they analyze the conformally Einstein product spaces, providing a classification for such spaces by using the idea of warped geometry. Our work, starting from a similar point of view, emphasizes the role of the conformal factor, laying out explicit formulas and the relative link to their geometric interpretation by using Penrose diagrams.

3. Conformal characterization

Firstly, we fix some important notation that we shall employ in this section. For the conformal factor Ω , we shall make explicit the dependence on variables, which for historically and interpretative reasons we identify with (t, x) .

Let us consider a generic 2-dimensional Lorentzian manifold (\mathcal{M}, g) , whose metric, thanks to Theorem 2.1, can be written as

$$g = \Omega(t, x) \cdot \eta \quad (1)$$

where $\Omega(t, x) > 0$ is a smooth function and η is the Minkowski metric defined in Def. 2.3. The Ricci scalar curvature R can be expressed as a function of Ω and its partial derivatives by

$$R = \frac{-(\partial_t \Omega(t, x))^2 + (\partial_x \Omega(t, x))^2 + \Omega(t, x) [\partial_t^2 \Omega(t, x) - \partial_x^2 \Omega(t, x)]}{(\Omega(t, x))^3}. \quad (2)$$

If we restrict ourselves to a traditional form for Ω , namely

$$\Omega(t, x) = e^{\omega(t, x)},$$

then Eq. (2) becomes:

$$R = (-\partial_t^2 \omega(t, x) + \partial_x^2 \omega(t, x)) e^{-\omega(t, x)}. \quad (3)$$

Remark 3.1. It is possible, by similar but much more cumbersome calculations, to compute the Ricci scalar curvature of a conformally flat manifold as a function of the conformal factor in a general dimension $n \geq 2$. Let us fix $\mathbf{x} = (x_1, \dots, x_{n-1})$ as the space coordinates, and t as the time coordinate. With this notation, we obtain the general formula

$$R = \left(\frac{1}{4}(n-1)(n-2) \left(\sum_{i=1}^{n-1} (\partial_{x_i} \omega(t, \mathbf{x}))^2 - (\partial_t \omega(t, \mathbf{x}))^2 \right) + (n-1) \left(\sum_{i=1}^{n-1} \partial_{x_i}^2 \omega(t, \mathbf{x}) - \partial_t^2 \omega(t, \mathbf{x}) \right) \right) e^{-\omega(t, \mathbf{x})}.$$

However, this expression is difficult to treat in its general form. As one may notice by direct computation, already for $n = 3$ the expression contains the square of the first order derivatives, and it requires a much deeper analysis:

$$R = \left(\frac{1}{2} \left(\sum_{i=1}^2 (\partial_{x_i} \omega(t, x_1, x_2))^2 - (\partial_t \omega(t, x_1, x_2))^2 \right) + 2 \left(\sum_{i=1}^2 \partial_{x_i}^2 \omega(t, x_1, x_2) - \partial_t^2 \omega(t, x_1, x_2) \right) \right) e^{-\omega(t, x_1, x_2)}.$$

We leave this, and higher dimensional cases, as outlook for future works. For the purpose of this article, we fix $n = 2$, in order to be able to provide a complete classification of 2-dimensional Lorentzian manifolds with Ricci scalar curvature, as we do in Theorem 2.1.

We remark an interesting property of the 2-dimensional Lorentzian manifolds.

Definition 3.1. *Let (\mathcal{M}, g) be a 2-dimensional Lorentzian manifold. We call it a Einstein manifold if the following condition holds:*

$$\text{Ric} = \kappa \cdot g, \quad (4)$$

for some constant $\kappa \in \mathbb{R}$, where Ric denotes the Ricci curvature tensor (here we are using the notation without indices as for the metric).

Proposition 3.1. *A 2-dimensional Lorentzian manifold is Einstein if and only if it has a constant Ricci scalar curvature R . Moreover the constant κ of Eq. (4) is equal to $\frac{R}{2}$.*

Proof. It is not difficult to compute the Ricci curvature tensor in terms of $\Omega = e^{\omega(t,x)}$. We get:

$$\begin{pmatrix} \frac{1}{2} (\Omega^{(0,2)}(t,x) - \Omega^{(2,0)}(t,x)) & 0 \\ 0 & \frac{1}{2} (\Omega^{(2,0)}(t,x) - \Omega^{(0,2)}(t,x)) \end{pmatrix}.$$

Now we see that the condition given by Eq. (4) holds if and only if

$$2\kappa e^{\omega(t,x)} = (-\partial_t^2 \omega(t,x) + \partial_x^2 \omega(t,x)).$$

Thanks to Eq. (3) we conclude the proof. \square

The conformal characterization of the Ricci scalar curvature given by Eq. (3) provide a very simple way to construct Lorentzian manifolds with constant curvature. The first step is to obtain the general form of a flat manifold. It is trivial how this is strictly connected with the classical change of variables as we shall see in Example 3.1.

Lemma 3.1. *The conformal factor $\Omega(t,x)$ of a generic 2-dimensional flat Lorentz manifold (i.e. with $R = 0$) can be written as*

$$\Omega(t,x) = \phi(x+t) \cdot \psi(x-t). \quad (5)$$

Proof. Let we fix $R = 0$. Thus, Eq. (3) leads to the one-dimensional wave equation. It is well known (see, e.g. case 8 of Sect. 4.1.3 in [15]) that its solutions can be expressed by the sum of a right traveling function $\hat{\phi}$ and a left traveling function $\hat{\psi}$. This means we can write:

$$\omega(t,x) = \hat{\phi}(x+t) + \hat{\psi}(x-t),$$

and so

$$\begin{aligned} \Omega(t,x) &= e^{\omega(t,x)} = \exp \left[\hat{\phi}(x+t) + \hat{\psi}(x-t) \right] = \\ &= \exp \left[\hat{\phi}(x+t) \right] \cdot \exp \left[\hat{\psi}(x-t) \right] = \phi(x+t) \cdot \psi(x-t). \end{aligned}$$

\square

Remark 3.2. *It is remarkable to notice that, setting $u = x+t$ and $v = x-t$, we have actually switched to the null coordinates. It is trivial to prove that, starting with the metric expressed in null coordinates, namely $ds^2 = dudv$, we get the same result of Lemma 3.1.*

Example 3.1. *Let we consider, for instance, the usual conformal transformation of the Minkowski space, which leads to the Penrose-Carter diagram (see, for instance, Chpt. 5 in [9]). It is easy to see, starting from the null coordinates¹, that the conformal factor could be written in terms of t and x as*


$$\Omega(t,x) = (\cos(t) + \cos(x))^{-2} = \frac{1}{4} \cos^{-2} \left(\frac{x+t}{2} \right) \cos^{-2} \left(\frac{x-t}{2} \right),$$

¹The trick is to set $\mu = \arctan(u)$ and $\nu = \arctan(v)$, with $-\frac{\pi}{2} \leq \mu \leq \frac{\pi}{2}$ and $-\frac{\pi}{2} \leq \nu \leq \frac{\pi}{2}$.

where now the range of t and x is given by:

$$-\pi < x < \pi \quad \text{and} \quad -\pi + |x| < t < \pi - |x|.$$

Thus, we recover, as expected, the form of Eq. (5).



(a) Minkowski space in classic coordinates.

(b) Penrose diagram of Minkowski space.

FIGURE 1. Here we provide a comparison between the Minkowski space in normal coordinates (a) and its conformal representation as Penrose-Carter diagram (b). The figures present the locus of point with the same *space interval* s^2 (also known in literature as *spacetime interval*), for several different values. We notice that the compactification in (b) produces the classical “diamond shape”, where the edges stand for the infinity.

Now we shift the focus on those conformal factors that depend only of one variable. We shall show (Example 3.2) how the general solution lead to a well known manifold: the de Sitter space.

Lemma 3.2. *Let us assume $R \neq 0$. The space-independent conformal factor $\Omega(t)$ of a generic 2-dimensional Lorentz manifold can be written as*

$$\Omega(t) = \frac{c_1 \left(-1 + \tanh^2 \left(\frac{1}{2} \sqrt{c_1(t+c_2)^2} \right) \right)}{2R}. \quad (6)$$

with c_1 and c_2 arbitrary constants.

Similarly, the time-independent conformal factor $\Omega(x)$ of a generic Lorentz manifold of dimension 2 can be written as

$$\Omega(x) = \frac{d_1 \left(1 - \tanh^2 \left(\frac{1}{2} \sqrt{d_1(x+d_2)^2} \right) \right)}{2R}, \quad (7)$$

with d_1 and d_2 arbitrary constants.

Proof. Here we just prove the space-independent case; the proof of the time-independent case is analogous. Assuming ω to be independent of x , Eq. (3) becomes

$$\frac{\partial^2}{\partial t^2}\omega(t) = R e^{\omega(t)}. \quad (8)$$

After some computations, we obtain the solutions to the second order nonlinear ODE (8):

$$\omega(t) = \log \left(\frac{c_1 \left(-1 + \tanh^2 \left(\frac{1}{2} \sqrt{c_1(c_1 + c_2)^2} \right) \right)}{2R} \right),$$

where $c_1, c_2 \in \mathbb{R}$. This concludes the proof. \square

Example 3.2. Let us consider Eq. (6). Setting $c_1 = -4$, $c_2 = 0$ and $R = 2$, we get:

$$\Omega(t) = 1 - \tanh^2(\sqrt{-t^2}) = \sec^2(t),$$

which, for the range

$$-\frac{\pi}{2} < t < \frac{\pi}{2} \quad \text{and} \quad 0 \leq x < 2\pi,$$

is exactly the conformal factor defining the 2-dimensional de Sitter space [7] in global coordinates (i.e. $ds^2 = \sec^2(t)(-dt^2 + dx^2)$).




FIGURE 2. Penrose-Carter diagram of the 2-dimensional de Sitter space in global coordinates.

Finally we have the general case, which allow us to construct several interesting manifolds and their Penrose- diagram as we present in Example 3.3.

Theorem 3.1. The general solution of Eq. (3) is given by

$$\omega(t, x) = \phi(x + t) + \psi(x - t) - 2 \log \left| k \int^{x+t} e^{\phi(\lambda)} d\lambda - \frac{R}{8k} \int^{x-t} e^{\psi(\lambda)} d\lambda + C \right|, \quad (9)$$

which, in the original Ω notation, becomes

$$\Omega(t, x) = \exp[\phi(x + t)] \cdot \exp[\psi(x - t)] \cdot \left(k \int^{x+t} e^{\phi(\lambda)} d\lambda - \frac{R}{8k} \int^{x-t} e^{\psi(\lambda)} d\lambda + C \right)^{-2}, \quad (10)$$

where ϕ and ψ arbitrary (smooth) functions, $k \neq 0$ and C is an arbitrary real constants. With $\int^x f(\lambda) d\lambda$ we denote the integration of $f(\lambda)$ with respect to λ and then the substitution $\lambda = x$.

In literature, equation (3) is called *modified* (or *generalized*) *Liouville equation*, and it is deeply studied in many works (for further details, see [11, 3, 15]).

Remark 3.3. As in Remark 3.2, we can rewrite Eq. (10) using null coordinates (u, v) (with $ds^2 = dudv$). So we get

$$\Omega(u, v) = e^{\phi(u)} \cdot e^{\psi(v)} \left(k \int^u e^{\phi(\lambda)} d\lambda - \frac{R}{8k} \int^v e^{\psi(\lambda)} d\lambda + C \right)^{-2} \quad (11)$$

with ϕ, ψ and C, k as in Theorem 3.1.




FIGURE 3. Example 3.3 lead us to the construction of the particular space (where the Ricci scalar curvature is $R = 2$) shown both in standard coordinates (a) and by the corresponding Penrose-Carter diagram (b).

Example 3.3. Let us consider Eq. (10) and let ϕ and ψ be the identity function. We also set $k = 1$, $R = 2$ and $C = 0$. So we obtain the following conformal factor in standard coordinates

$$\Omega_1(t, x) = e^{2x} \left(e^{x+t} - \frac{1}{4} e^{x-t} \right)^{-2},$$

which defines a metric of scalar curvature $R = 2$; see Figure 3(a) for a visualization. By using Remark 3.3 and Eq. (11), it is possible to provide the Penrose-diagram (see Example 3.1) as one can see in Figure 3. In this case the conformal factor is given by:

$$\Omega_2(t, x) = \frac{\exp(\tan(\frac{t+x}{2})) \exp(\tan(\frac{x-t}{2}))}{(2 \cos(\frac{t+x}{2}) \cos(\frac{x-t}{2}))^2 (\exp(\tan(\frac{t+x}{2})) - \frac{1}{4} \exp(\tan(\frac{x-t}{2})))^2},$$

where the range of t and x is:

$$-\pi < x < \pi \quad \text{and} \quad -\pi + |x| < t < \pi - |x|.$$

See Figure 3(b) for a useful visualization.

Remark 3.4. It is now possible, by varying parameters, to obtain all the possible Lorentzian space with constant Ricci scalar curvature and applying the same argument we used in this work, also their corresponding Penrose-Carter diagram.

4. Conclusions

In this note, we provided a constructive way to develop a conformal characterization of 2-dimensional Lorentzian manifolds with constant Ricci scalar curvature. The tools we used are very classical, but their application allow us to create a suggestive set of Lorentzian manifolds, starting from the conformal factor which, as we have shown, identifies them. Many useful examples have been proposed to explain the power of such construction. In the near future, we would like to investigate other similar applications using the same approach to characterize 2-dimensional Lorentzian manifolds with a general (i. e. not constant) Ricci scalar curvature, or for the 3-dimensional Einstein manifolds (with the respective geometric interpretation).

Acknowledgments: The authors would like to thank Daniele Taufer for the fruitful exchanges during the writing of this paper. The authors would also like to acknowledge the insightful feedback given by the reviewers. NC and MS would like to thank the University of Pavia and the University of Trento, respectively, for supporting their research.

References

- [1] J. K. Beem. *Global Lorentzian Geometry*. Chapman & Hall/CRC Pure and Applied Mathematics. CRC Press, 2017.
- [2] A. L. Besse. *Einstein manifolds*. Springer Science & Business Media, 2007.
- [3] R. K. Bullough and P. J. Caudrey, editors. *Solitons*. Springer-Verlag, 1980.
- [4] J. Burgos. Two-dimensional Minkowski causal automorphisms and conformal maps. *Class. Quantum Gravity*, 30, 05 2013.
- [5] B.-Y. Chen. *Pseudo-Riemannian Geometry, δ -Invariants and Applications*. World Scientific, 2011.
- [6] P. Collas. General relativity in two- and three-dimensional space-times. *Am. J. Phys.*, 45:833–837, 09 1977.
- [7] H. S. M. Coxeter. A geometrical background for de Sitter’s world. *Am. Math. Mon.*, 50(4):217–228, 1943.
- [8] D. Grumiller, W. Kummer, and D.V. Vassilevich. Dilaton gravity in two-dimensions. *Phys. Rept.*, 369:327–430, 2002.
- [9] S. Hawking and G. Ellis. *The Large Scale Structure of Space-Time*. Cambridge University Press, 1973.
- [10] W. Kühnel and H.-B. Rademacher. Conformally Einstein product spaces. *Differential Geometry and its Applications*, 49:65 – 96, 2016.
- [11] J. Liouville. Sur l’équation aux différences partielles: $\partial^2 \log \lambda / \partial u \partial v \pm \lambda / 2a^2 = 0$. *Journal de Mathématiques Pures et Appliquées*, 18:71–72, 1853.
- [12] C. W. Norman. *Undergraduate algebra: a first course*. Oxford University Press, USA, 1986.
- [13] K. Papadopoulos, N. Kurt, and B. Papadopoulos. On the causal and topological structure of the 2-dimensional minkowski space. *Universe*, 02 2019.
- [14] F. Pfäffle. *Quantum Field Theory on Curved Spacetimes - Lecture Notes in Physics (Vol. 786)*, chapter Lorentzian Manifolds. Springer, Berlin, Heidelberg, 2009.
- [15] A. D. Polyanin and V. F. Zaitsev. *Handbook of nonlinear partial differential equations*. CRC press, 2004.
- [16] M. Schottenloher. *A mathematical introduction to conformal field theory*, volume 759. Springer, 2008.
- [17] I. Terek and A. Lymberopoulos. A local characterization for constant curvature metrics in 2-dimensional lorentz manifolds. *preprint, arXiv:1605.07573*, 2016.
- [18] R. M. Wald. *General Relativity*. Chicago Univ. Press, Chicago, IL, 1984.
- [19] R. M. Wald and U. Yurtsever. General proof of the averaged null energy condition for a massless scalar field in two-dimensional curved space-time. *Phys. Rev. D*, 44:403–416, 1991.