

## LIE SYMMETRY ANALYSIS AND CONSERVATION LAWS FOR TIME FRACTIONAL COUPLED WHITHAM-BROER- KAUP EQUATIONS

Hadi Roohani GHEHSAREH<sup>1</sup>, Ahmad MAJLESI<sup>2</sup>, Ali ZAGHIAN<sup>3</sup>

*In the current work, time fractional coupled Whitham-Broer-Kaup equations which describes the anomalous bidirectional propagation of long waves in shallow water is investigated. A Lie symmetry analysis is formulated and used to the governing model. The symmetry reductions of the model are constructed and the system of nonlinear time fractional partial differential equations is similarity reduced to a system of nonlinear fractional ordinary differential equations in Erdelyi-Kober derivative sense. Moreover, the resultant symmetry generators are used to calculate conserved vectors for the time fractional problem. Two different kinds of conservation laws of the problem have been constructed.*

**Keywords:** Lie symmetry analysis; Time fractional coupled Whitham-Broer-Kaup equations; Erdelyi-Kober operators; Conservation laws

### 1. Introduction

In recent decades theory of fractional calculus [1, 2] has gained notable attention of many researchers in science and engineering due to its high ability for describing various complicated natural phenomena. Many of anomalous phenomena and complex processes in natural science have been successfully described by using the theory of fractional calculus and mathematically modeled as fractional differential or integral equations[3, 4, 5, 6]. The main advantage of fractional modeling is that they are excellent tools to appropriately characterize hereditary and inherent memory properties of the anomalous phenomena. In recent years several semi-analytical methods and numerical techniques have been developed and employed for solving and investigating various practical problems which have been mathematically modeled as fractional equations [7, 8, 9, 10, 11, 12, 13, 14, 15, 16].

In this study the following important mathematical model, the time fractional Whitham-Broer-Kaup equations, which describes the anomalous bidirectional propagation of long waves in shallow water will be investigated[17].

<sup>1</sup> Prof., Dept. of Mathematics, Malek Ashtar University Of Technology, Shahin Shahr , 34149-16818, Iran, e-mail: hadiroohani61@gmail.com

<sup>2</sup> PhD student, Dept.of Mathematics, Malek Ashtar University Of Technology, Shahin Shahr , 34149-16818, Iran, e-mail: ahmad.majlessi@gmail.com

<sup>3</sup> Prof., Dept.of Mathematics, Malek Ashtar University Of Technology, Shahin Shahr , 34149-16818, Iran, e-mail: ali\_zaghian1338@yahoo.com

$$\begin{aligned} \frac{\partial^\alpha u}{\partial t^\alpha} + \frac{\partial v}{\partial x} + u \frac{\partial u}{\partial x} + q \frac{\partial^2 u}{\partial x^2} &= 0, \\ \frac{\partial^\alpha v}{\partial t^\alpha} + \frac{\partial(uv)}{\partial x} + p \frac{\partial^3 u}{\partial x^3} - q \frac{\partial^2 v}{\partial x^2} &= 0, \end{aligned} \quad (1)$$

where  $u(x, t)$  and  $v(x, t)$  denote the fluid velocity along the horizontal direction and vertical displacement of the fluid from its equilibrium position, respectively.

The constants  $p$  and  $q$  ( $(p, q) \neq (0, 0)$ ) are diffusion coefficients and  $\frac{\partial^\alpha}{\partial t^\alpha}(.)$  denotes the Riemann-Liouville partial derivative operator of order  $\alpha$  ( $0 < \alpha \leq 1$ ) with respect to time component,  $t$ , which is defined as follows :

$$\frac{\partial^\alpha}{\partial t^\alpha}(u(x, t)) = \begin{cases} \frac{1}{\Gamma(n-\alpha)} \frac{\partial^n}{\partial t^n} \int_0^t (t-\tau)^{m-\alpha-1} u(x, \tau) d\tau, & m-1 < \alpha < m, m \in \mathbb{N}, \\ \frac{\partial^n}{\partial t^n}(u(x, t)), & \alpha = n \in \mathbb{N}, \end{cases}$$

where  $\Gamma(.)$  is the gamma function. For  $\alpha = 1$  the anomalous bidirectional propagation model (1) is reduced to the classical Whitham-Broer-Kaup problem. Several numerical and semi-analytical methods have been formulated and employed to investigate the classical Whitham-Broer-Kaup problem[18, 19, 20, 21, 22]. The classical Whitham-Broer-Kaup (WBK) equation has been widely used to study of solitary wave theory in shallow water. However, the classical Whitham-Broer-Kaup (WBK) equation is not adequate to describe complicated mechanism of propagation of shallow water waves in porous medium, such as tsunami wave propagation. In these situations the temporal derivative in the WBK equation and corresponds to the variation in the flux,  $\frac{\partial}{\partial t}$ , should be replaced

with fractional time derivative,  $\frac{\partial^\alpha}{\partial t^\alpha}$ , which properly describes variation in the flux through the fractal boundary, where  $\alpha$  denotes the fractal dimensions of the porous medium [23]. Saha Ray [17] has discussed the time-fractional WBK equations (1) and approximated the traveling wave solutions of the model in their generalized Taylor expansion forms. In [24] a semi-analytical technique, called the residual power series method is formulated and used to approximate traveling solutions of the model (1). By choosing  $p = 1$  and  $q = 0$  in the (1) the so-called time fractional modified Boussinesq system is concluded. While if  $p = 0$  and

$q \neq 0$  another special case of the time fractional coupled Whitham-Broer-Kaup equations, called time fractional coupled approximate long wave equations is obtained. In [25] the time-space fractional Whitham-Broer-Kaup equations has been investigated. Recently, Amjad Ali et.al employed a numerical method based on the Adomian decomposition method coupled with the Laplace transforms to calculate the approximate traveling wave solutions of the time-space fractional coupled Whitham-Broer-Kaup equations[26].

In recent years, Lie symmetry analysis has been developed and widely used to deal with several types of complicated nonlinear differential problems[27, 28, 29, 30]. Buckwar and Luchko in [31] computed invariant solutions of fractional differential equations by employing the scaling transformations. Moreover, a symmetry group of scaling transformations for time-space fractional partial differential equation is extracted in [32]. Gazizov et al. introduced a prolongation formula for the Riemann-Liouville fractional derivative operator[33]. In [34] they utilized the proposed Lie point symmetry method for solving the nonlinear time fractional diffusion problem. Recently, Hashemi et al. formulated and used the Lie symmetry approach for investigating various types of the fractional differential equations[35,36,37,38,39,40,41]. Singla and Gupta extracted a Lie point symmetry analysis for systems of coupled time fractional partial differential equations [42,43]. Moreover, they developed the Lie symmetry approach for space-time fractional systems of partial differential equations [44, 45]. In this study, the expanded Lie symmetry approach proposed by Singla and Gupta is utilized for symmetry analysis and similarity reductions of the time fractional Whitham-Broer-Kaup equations (1).

The idea of conservation law plays an important role to analyze the fundamental properties of the physical models[46]. Relation between symmetries of differential equations and conservation laws is explained by the Noether's theorem [47]. The Noether's theorem is valid for differential equations having Lagrangians. The classical Noether's theorem have been generalized and employed to find conservation laws for several fractional differential equations having fractional Lagrangians [48,49,50,51]. Ibragimov in [52] extracted a new generalized conservation theorem based on the adjoint equations for the nonlinear differential equations not having Lagrangians. Lukashchuk has used the new conservation theorem to find conservation laws for time fractional subdiffusion and diffusion-wave equations[53]. Gazizov et al. found conservation laws for the time-fractional Kompaneets equations based on the generalization of fractional Noether's operator [54]. Very recently, Singla and Gupta in [55,56] have extracted the fractional Noether's operators to calculate conserved vectors of the time and space-time fractional nonlinear systems of partial differential equations, respectively. Majlesi et.al. performed a Lie symmetry analysis on a coupled system of time fractional Jaulent-Miodek equations and constructed the

conservation laws of the problem [57]. In the current work the generalization of fractional Noether's operator proposed by Singla and Gupta [55] is employed for calculating conserved vectors for the governing system of nonlinear time fractional partial differential equation (1).

## 2. Lie symmetry analysis and similarity reductions of the time fractional Whitham-Broer-Kaup system

In this section an invariant analysis for the time fractional Whitham-Broer-Kaup system (1) would be presented. Moreover, the symmetry reductions of the governing time fractional nonlinear system based on the symmetry groups are investigated. For this purpose, firstly the main details of Lie symmetry analysis for systems of time fractional PDEs are briefly described.

### 2.1 Description of the Lie symmetry analysis for systems of time fractional PDEs

Here we consider a coupled system of two time fractional nonlinear PDEs as follows[42]:

$$\Delta_1 \equiv \frac{\partial^\alpha u}{\partial t^\alpha} - F(x, t, u, v, u_x, v_x, \dots) = 0, \quad (2)$$

$$\Delta_2 \equiv \frac{\partial^\alpha v}{\partial t^\alpha} - G(x, t, u, v, u_x, v_x, \dots) = 0,$$

where  $\{u(x, t), v(x, t)\}$  and  $\{x, t\}$  are dependent and independent variables respectively.  $\alpha > 0$  is a real number and  $\frac{\partial^\alpha}{\partial t^\alpha}(\cdot)$  denotes the Riemann-Liouville partial derivative operator and subscripts denote integer partial derivatives. According to the Lie symmetry analysis, assume the time fractional system (2) is invariant under the following one parameter Lie group of transformations :

$$\begin{aligned} \bar{x} &= x + \varepsilon \xi(x, t, u, v) + O(\varepsilon^2), \\ \bar{t} &= t + \varepsilon \tau(x, t, u, v) + O(\varepsilon^2), \\ \bar{u} &= u + \varepsilon \eta(x, t, u, v) + O(\varepsilon^2), \\ \bar{v} &= v + \varepsilon \phi(x, t, u, v) + O(\varepsilon^2), \\ \frac{\partial^\alpha \bar{u}}{\partial \bar{t}^\alpha} &= \frac{\partial^\alpha u}{\partial t^\alpha} + \varepsilon \eta^{\alpha, t} + O(\varepsilon^2), \\ \frac{\partial^\alpha \bar{v}}{\partial \bar{t}^\alpha} &= \frac{\partial^\alpha v}{\partial t^\alpha} + \varepsilon \phi^{\alpha, t} + O(\varepsilon^2), \end{aligned} \quad (3)$$

$$\begin{aligned}\frac{\partial^j \bar{u}}{\partial \bar{x}^j} &= \frac{\partial^j u}{\partial x^j} + \varepsilon \eta^{j,x} + O(\varepsilon^2), \quad j = 1, 2, 3, \\ \frac{\partial^j \bar{v}}{\partial \bar{x}^j} &= \frac{\partial^j v}{\partial x^j} + \varepsilon \phi^{j,x} + O(\varepsilon^2), \quad j = 1, 2, 3,\end{aligned}$$

where  $\varepsilon$  is a parameter,  $(\xi, \tau, \eta, \phi)$  is the set of infinitesimals and  $\eta^{\alpha,t}, \phi^{\alpha,t}$  and  $\eta^{j,x}, \phi^{j,x}, (j = 1, 2,)$  are extended infinitesimals of order  $\alpha$  and  $j$  respectively. The infinitesimal generator of the time fractional system (2) is a vector field as follows:

$$V = \xi(x, t, u, v) \frac{\partial}{\partial x} + \tau(x, t, u, v) \frac{\partial}{\partial t} + \eta(x, t, u, v) \frac{\partial}{\partial u} + \phi(x, t, u, v) \frac{\partial}{\partial v}. \quad (4)$$

The vector field (4) generates a symmetry of (2) provided that it admits the following infinitesimal invariance criteria:

$$\begin{aligned}Pr^\alpha V(\Delta_1)|_{\Delta_1=0, \Delta_2=0} &= 0, \\ Pr^\alpha V(\Delta_2)|_{\Delta_1=0, \Delta_2=0} &= 0,\end{aligned} \quad (5)$$

where  $Pr^\alpha V$  is the generalized fractional prolongation operator[42,43] :

$$Pr^{\alpha, m, n} V = V + \eta^{\alpha,t} \frac{\partial}{\partial u_t^{(\alpha)}} + \eta^{1,x} \frac{\partial}{\partial u_x} + \eta^{m,x} \frac{\partial}{\partial u_{mx}} + \phi^{\alpha,t} \frac{\partial}{\partial v_t^{(\alpha)}} + \phi^{1,x} \frac{\partial}{\partial v_x} + \phi^{n,x} \frac{\partial}{\partial v_{nx}}, \quad (6)$$

where  $m, n \in \mathbb{N}$  denote orders of partial differential equations (2). In the above relation the integer order extended infinitesimals  $\eta^{j,x}$  and  $\phi^{j,x}$  are defined as follows:

$$\begin{aligned}\eta^{j,x} &= D_x \eta^{j-1,x} - (D_x \xi) \frac{\partial^j u}{\partial x^j} - (D_x \tau) \frac{\partial}{\partial t} \left( \frac{\partial^{j-1} u}{\partial x^{j-1}} \right), \\ \phi^{j,x} &= D_x \phi^{j-1,x} - (D_x \xi) \frac{\partial^j v}{\partial x^j} - (D_x \tau) \frac{\partial}{\partial t} \left( \frac{\partial^{j-1} v}{\partial x^{j-1}} \right), \quad j = 1, 2,,\end{aligned}$$

where  $D_x$  denotes the total derivative operator with respect to  $x$

$$D_x = \frac{\partial}{\partial x} + u_x \frac{\partial}{\partial u} + u_{xx} \frac{\partial}{\partial u_x} + v_x \frac{\partial}{\partial v} + v_{xx} \frac{\partial}{\partial v_x} + \dots,$$

Moreover the  $\alpha$ -order extended infinitesimal  $\eta^{\alpha,t}$  is defined as follows:

$$\eta^{\alpha,t} = D_t^\alpha \eta + \xi D_t^\alpha (u_x) - D_t^\alpha (\xi u_x) + D_t^\alpha (D_t(\tau)u) - D_t^{\alpha+1} (\tau u) + \tau D_t^{\alpha+1} (u),$$

where  $D_t^\alpha$  is the total fractional derivative with respect to  $t$ . Using the generalized Leibniz rule [2] and generalized chain rule [5] and after some simplifications the  $\alpha$ -order extended infinitesimal  $\eta^{\alpha,t}$  is given in the following form (see for more details [42, 43]) :

$$\begin{aligned}
\eta^{\alpha,t} = & \frac{\partial^{\alpha\eta}}{\partial t^\alpha} + (\eta_u - \alpha D_t(\tau)) \frac{\partial^\alpha u}{\partial t^\alpha} - u \frac{\partial^\alpha \eta_u}{\partial t^\alpha} + (\eta_v \frac{\partial^\alpha v}{\partial t^\alpha} - v \frac{\partial^\alpha \eta_v}{\partial t^\alpha}) \\
& + \sum_{i=1}^{\infty} \left[ \binom{\alpha}{i} \frac{\partial^i \eta_u}{\partial t^i} - \binom{\alpha}{i+1} D_t^{i+1}(\tau) \right] D_t^{\alpha-i}(u) + \sum_{i=1}^{\infty} \left[ \binom{\alpha}{i} \frac{\partial^i \eta_v}{\partial t^i} D_t^{\alpha-i}(v) \right] \\
& - \sum_{i=1}^{\infty} \left[ \binom{\alpha}{i} D_t^i(\xi) D_t^{\alpha-i}(u_x) + \zeta_1 + \zeta_2 \right], \tag{7}
\end{aligned}$$

where  $D_t$  denotes the total derivative operator with respect to  $t$  and

$$\zeta_1 = \sum_{i=2}^{\infty} \sum_{j=2}^i \sum_{r=2}^{j-1} \sum_{s=0}^{r-1} \binom{\alpha}{i} \binom{i}{j} \binom{r}{s} \frac{t^{i-\alpha}}{r! \Gamma(i-\alpha+1)} (-1)^s u^s \frac{\partial^j (u^{r-s})}{\partial t^j} \frac{\partial^{i-j+r} \eta}{\partial t^{i-j} \partial u^r},$$

$$\zeta_2 = \sum_{i=2}^{\infty} \sum_{j=2}^i \sum_{r=2}^{j-1} \sum_{s=0}^{r-1} \binom{\alpha}{i} \binom{i}{j} \binom{r}{s} \frac{t^{i-\alpha}}{r! \Gamma(i-\alpha+1)} (-1)^s v^s \frac{\partial^j (v^{r-s})}{\partial t^j} \frac{\partial^{i-j+r} \eta}{\partial t^{i-j} \partial v^r}.$$

Similarity, the  $\alpha$ -order extended infinitesimal  $\phi^{\alpha,t}$  can be simplified as follows :

$$\begin{aligned}
\phi^{\alpha,t} = & \frac{\partial^{\alpha\phi}}{\partial t^\alpha} + (\phi_v - \alpha D_t(\tau)) \frac{\partial^\alpha v}{\partial t^\alpha} - v \frac{\partial^\alpha \phi_v}{\partial t^\alpha} + (\phi_u \frac{\partial^\alpha u}{\partial t^\alpha} - u \frac{\partial^\alpha \phi_u}{\partial t^\alpha}) \\
& + \sum_{i=1}^{\infty} \left[ \binom{\alpha}{i} \frac{\partial^i \phi_v}{\partial t^i} - \binom{\alpha}{i+1} D_t^{i+1}(\tau) \right] D_t^{\alpha-i}(v) + \sum_{i=1}^{\infty} \left[ \binom{\alpha}{i} \frac{\partial^i \phi_u}{\partial t^i} D_t^{\alpha-i}(u) \right] \\
& - \sum_{i=1}^{\infty} \left[ \binom{\alpha}{i} D_t^i(\xi) D_t^{\alpha-i}(v_x) + \rho_1 + \rho_2 \right], \tag{8}
\end{aligned}$$

where

$$\rho_1 = \sum_{i=2}^{\infty} \sum_{j=2}^i \sum_{r=2}^{j-1} \sum_{s=0}^{r-1} \binom{\alpha}{i} \binom{i}{j} \binom{r}{s} \frac{t^{i-\alpha}}{r! \Gamma(i-\alpha+1)} (-1)^s u^s \frac{\partial^j (u^{r-s})}{\partial t^j} \frac{\partial^{i-j+r} \phi}{\partial t^{i-j} \partial u^r},$$

$$\rho_2 = \sum_{i=2}^{\infty} \sum_{j=2}^i \sum_{r=2}^{j-1} \sum_{s=0}^{r-1} \binom{\alpha}{i} \binom{i}{j} \binom{r}{s} \frac{t^{i-\alpha}}{r! \Gamma(i-\alpha+1)} (-1)^s v^s \frac{\partial^j (v^{r-s})}{\partial t^j} \frac{\partial^{i-j+r} \phi}{\partial t^{i-j} \partial v^r},$$

From the above relations it can be easily concluded that whenever infinitesimals  $\phi$  and  $\eta$  are linear with respect to each dependent variables  $u$  and  $v$ , the expressions  $\zeta_1, \zeta_2, \rho_1$  and  $\rho_2$  vanish identically.

## 2.2 The symmetry reductions of the time fractional Whitham-Broer-Kaup system

Here, the proposed Lie symmetry analysis is formulated and employed to reduce the problem (1). For this purpose, by imposing the prolongations (6)  $Pr^{\alpha,2,1}V(\Delta_1)$  and  $Pr^{\alpha,3,2}V(\Delta_2)$  to problem (1) under group of transformations (3), the following invariance criterions are concluded:

$$\begin{aligned} & [\eta^{\alpha,t} + \phi^{1,x} + \eta u_x + u \eta^{1,x} + q \eta^{2,x}]_{\Delta_1=0, \Delta_2=0} = 0, \\ & [\phi^{\alpha,t} + p \eta^{3,x} + u \phi^{1,x} + v \eta^{1,x} + \eta v_x + \phi u_x - q \phi^{2,x}]_{\Delta_1=0, \Delta_2=0} = 0. \end{aligned} \quad (9)$$

By substituting the integer and  $\alpha$  order extended infinitesimals into (9) and equating the coefficients of various powers and partial derivatives of independent variables  $u$  and  $v$  to zero, a set of fractional and classical partial differential equations with respect to the variables  $\xi, \tau, \eta$  and  $\phi$  is obtained. By solving the resultant system of differential equations symbolically the following values for infinitesimals functions are computed:

$$\xi = c_1 + c_2 \alpha x, \quad \tau = c_3 + 2t c_2, \quad \eta = -\alpha c_2 u, \quad \phi = -2\alpha c_2 v, \quad (10)$$

where  $c_1, c_2$  and  $c_3$  are free constant parameters. Notice that the lower limit of the Riemann-Liouville partial derivative operator is fixed, so to ensure that it is invariant under group of transformations (3), the following initial condition should be satisfied.

$$\tau(x, t, u, v)|_{t=0} = 0. \quad (11)$$

So  $c_3 = 0$  and the following sets of the infinitesimal generators are obtained:

$$V_1 = \alpha x \frac{\partial}{\partial x} + 2t \frac{\partial}{\partial t} - \alpha u \frac{\partial}{\partial u} - 2\alpha v \frac{\partial}{\partial v}, \quad V_2 = \frac{\partial}{\partial x}. \quad (12)$$

Clearly, the above vector fields form a closed Lie algebra:

$$[V_1, V_1] = [V_2, V_1] = 0, \quad [V_1, V_2] = -\alpha V_2 = -[V_2, V_1].$$

For the infinitesimal generator  $V_1$ , the following characteristic equations is concluded :

$$\frac{dx}{\alpha x} = \frac{dt}{2t} = \frac{du}{-\alpha u} = \frac{dv}{-2\alpha v}.$$

By solving the above characteristic equations the following similarity variable and invariant solutions are obtained.

$$z = xt^{-\frac{\alpha}{2}}, \quad u(x, t) = t^{\frac{\alpha}{2}} F(z), \quad v(x, t) = t^{-\alpha} G(z), \quad (13)$$

where  $z$  and  $F(z), G(z)$  are new independent variable and dependent variables respectively.

Now, the reduction form of the time fractional system (1) with respect to the above presented symmetry generators is given by the following theorem.

**Theorem 1.** Corresponding to the infinitesimal generator  $V_1$ , the similarity variable  $z = xt^{-\frac{\alpha}{2}}$  with the similarity transformations  $u(x, t) = t^{-\frac{\alpha}{2}}F(z)$  and  $v(x, t) = t^{-\alpha}G(z)$ , reduced the time fractional system of partial differential equations (1) to the following system of fractional ordinary differential of equations:

$$\begin{aligned} (P_{\frac{2}{\alpha}}^{1-\frac{3\alpha}{2}}F)(z) &= -G'(z) + F(z)F'(z) + qF''(z), \\ (P_{\frac{2}{\alpha}}^{1-\frac{3\alpha}{2}}G)(z) &= -pF''(z) + (F(z)G(z))' - qG''(z), \end{aligned} \quad (14)$$

where the left-hand side Erdelyi-Kober fractional differential operator,  $(P_{\zeta}^{\rho, \alpha})$ , is defined as[58] :

$$(P_{\zeta}^{\rho, \alpha}H)(z) = \prod_{j=0}^{m-1} (\rho + j - \frac{1}{\zeta}z \frac{d}{dz})(K_{\zeta}^{\rho+\alpha, m-\alpha}H)(z), \quad z > 0, \alpha > 0, \quad (15)$$

where

$$m = \begin{cases} [\alpha] + 1 & \text{if } \alpha \notin \mathbb{N}, \\ \alpha & \text{if } \alpha \in \mathbb{N}, \end{cases}$$

and

$$(K_{\zeta}^{\rho, \alpha}H)(z) = \begin{cases} \frac{1}{\Gamma(\alpha)} \int_1^{\infty} (\iota - 1)^{\alpha-1} \iota^{-(\rho+\alpha)} H(z \iota^{\frac{1}{\zeta}}) d\iota & \text{if } \alpha > 0 \\ H(z) & \text{if } \alpha = 0 \end{cases}$$

**Proof:** According to the similarity solutions (13), for  $n-1 < \alpha < n$  ( $n \in \mathbb{N}$ ) the  $\alpha$ -order Riemann-Liouville time fractional derivative for  $u(x, t)$  is given as follows:

$$\frac{\partial^{\alpha}u}{\partial t^{\alpha}} = \frac{1}{\Gamma(n-\alpha)} \frac{\partial^n}{\partial t^n} \int_0^t (t-s)^{n-\alpha-1} s^{-\frac{\alpha}{2}} F(xs^{-\frac{\alpha}{2}}) ds, \quad (16)$$

Now by using the change of variable,  $\iota = \frac{t}{s}$ , the relation (16) is transformed to following form :

$$\frac{\partial^{\alpha}u}{\partial t^{\alpha}} = \frac{\partial^n}{\partial t^n} \left[ \frac{t^{n-\frac{3\alpha}{2}}}{\Gamma(n-\alpha)} \int_1^{\infty} (\iota - 1)^{n-\alpha-1} \iota^{-(n-\frac{3\alpha}{2}+1)} F(z \iota^{\frac{\alpha}{2}}) d\iota \right], \quad (17)$$

by comparing with the Erdelyi-Kober fractional differential operator, the above relation can be presented as follows:

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \frac{\partial^n}{\partial t^n} [t^{n-\frac{3}{2}\alpha} (K_{\frac{2}{\alpha}}^{1-\frac{\alpha}{2}, n-\alpha} F)(z)]. \quad (18)$$

So we have,

$$\begin{aligned} \frac{\partial^\alpha u}{\partial t^\alpha} &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left( \frac{\partial}{\partial t} [t^{n-\frac{3}{2}\alpha} (K_{\frac{2}{\alpha}}^{1-\frac{\alpha}{2}, n-\alpha} F)(z)] \right) \\ &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[ (n - \frac{3}{2}\alpha) t^{n-\frac{3}{2}\alpha-1} (K_{\frac{2}{\alpha}}^{1-\frac{\alpha}{2}, n-\alpha} F)(z) + t^{n-\frac{3}{2}\alpha} \frac{d}{dz} (K_{\frac{2}{\alpha}}^{1-\frac{\alpha}{2}, n-\alpha} F)(z) \frac{\partial z}{\partial t} \right] \\ &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[ t^{n-\frac{3}{2}\alpha-1} (n - \frac{3}{2}\alpha + t \frac{\partial z}{\partial t} \frac{d}{dz}) (K_{\frac{2}{\alpha}}^{1-\frac{\alpha}{2}, n-\alpha} F)(z) \right] \\ &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[ t^{n-\frac{3}{2}\alpha-1} (n - \frac{3}{2}\alpha - \frac{\alpha}{2} z \frac{d}{dz}) (K_{\frac{2}{\alpha}}^{1-\frac{\alpha}{2}, n-\alpha} F)(z) \right]. \end{aligned} \quad (19)$$

Following the above process for  $(n-1)$  times, we easily obtain:

$$\begin{aligned} \frac{\partial^\alpha u}{\partial t^\alpha} &= \frac{\partial^n}{\partial t^n} [t^{n-\frac{3}{2}\alpha} (K_{\frac{2}{\alpha}}^{1-\frac{\alpha}{2}, n-\alpha} F)(z)] \\ &= t^{-\frac{3}{2}\alpha} \prod_{j=0}^{n-1} \left( 1 - \frac{3}{2}\alpha + j - \frac{\alpha}{2} z \frac{d}{dz} \right) (K_{\frac{2}{\alpha}}^{1-\frac{\alpha}{2}, n-\alpha} F)(z) \\ &= t^{-\frac{3}{2}\alpha} (P_{\frac{2}{\alpha}}^{1-\frac{3\alpha}{2}, \alpha} F)(z). \end{aligned} \quad (20)$$

In similar manner, the  $\alpha$ -order Riemann-Liouville time fractional derivative for  $v(x, t)$  with respect to the similarity transforms (13) can be expressed as follows:

$$\frac{\partial^\alpha v}{\partial t^\alpha} = t^{-2\alpha} (P_{\frac{2}{\alpha}}^{1-2\alpha, \alpha} G)(z). \quad (21)$$

Moreover for  $\alpha = n \in \mathbb{N}$  we have:

$$\begin{aligned} \frac{\partial^\alpha u}{\partial t^\alpha} &= \frac{\partial^n u}{\partial t^n} = \frac{\partial^n}{\partial t^n} (t^{-\frac{n}{2}} F(z)) \\ &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[ \frac{\partial}{\partial t} (t^{-\frac{n}{2}} F(z)) \right] \\ &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[ t^{-\frac{n-1}{2}} \left( -\frac{n}{2} - \frac{n}{2} z \frac{d}{dz} \right) F(z) \right] \\ &\vdots \\ &= t^{-\frac{3}{2}n} \prod_{j=0}^{n-1} \left( 1 - \frac{3}{2}n + j - \frac{n}{2} z \frac{d}{dz} \right) F(z) \end{aligned}$$

$$= t^{-\frac{3}{2}n} (P_{\frac{2}{n}}^{1-\frac{3n}{2}, n} F)(z),$$

and similarly, for  $\alpha = n \in \mathbb{N}$  we have:

$$\frac{\partial^\alpha v}{\partial t^\alpha} = \frac{\partial^n v}{\partial t^n} = t^{-2n} (P_{\frac{2}{n}}^{1-2n, n} G)(z).$$

Now by substituting the proposed similarity transformations (13) in the main problem (1) and from relations (20) and (21) the reduced system of fractional ordinary equations (14) is concluded.

Moreover, for the infinitesimal generator  $V_2$ , by solving the associated characteristic equations the invariant solutions of the problem (1) are obtained as  $u(x, t) = f(t)$  and  $v(x, t) = g(t)$ . Substituting the above obtained group of invariant solutions in the governing problem (1), the following reduced system is concluded:

$$\begin{cases} \frac{\partial^\alpha f(t)}{\partial t^\alpha} = 0 \\ \frac{\partial^\alpha g(t)}{\partial t^\alpha} = 0 \end{cases}$$

Integrating both sides of the above system, the following invariant solutions are arrived:

$$f(t) = \frac{\kappa_1}{\Gamma(\alpha)} t^{\alpha-1}, \quad g(t) = \frac{\kappa_2}{\Gamma(\alpha)} t^{\alpha-1}, \quad (22)$$

where  $\kappa_1$  and  $\kappa_2$  are arbitrary constants.

### 3. Conservation laws of the time fractional Whitham-Broer-Kaup system

In this section a new approach which has been developed and implemented to the time-fractional system of PDEs [55] is employed to construct conservation laws for the time fractional Whitham-Broer-Kaup system (1). According to the approach proposed by Singla and Gupta[55], the formal Lagrangian for the time fractional PDEs system (1) is considered as follows:

$$\begin{aligned} L = & f(x, t) \left[ \frac{\partial^\alpha u}{\partial t^\alpha} + v_x(x, t) + u(x, t)u_x(x, t) + qu_{xx}(x, t) \right] \\ & + g(x, t) \left[ \frac{\partial^\alpha v}{\partial t^\alpha} + pu_{xxx}(x, t) + u(x, t)v_x(x, t) + u_x(x, t)v(x, t) - qv_{xx}(x, t) \right], \end{aligned} \quad (23)$$

where  $f(x, t)$  and  $g(x, t)$  are newly introduced dependent variables. The adjoint equations for the formal Lagrangian operator (23) are calculated as follows:

$$\frac{\delta L}{\delta u} = (D_t^\alpha)^* f - u(x, t) f_x(x, t) - v(x, t) g_x(x, t) + q f_{xx}(x, t) - p g_{xxx}(x, t) = 0, \quad (24)$$

and

$$\frac{\delta L}{\delta v} = (D_t^\alpha)^* g - f_x(x, t) - g_x(x, t) u(x, t) - q g_{xx}(x, t) = 0, \quad (25)$$

where the Euler-Lagrange operator,  $\frac{\delta}{\delta \omega}$ , is defined as :

$$\frac{\delta}{\delta \omega} = \frac{\partial}{\partial \omega} + (D_t^\alpha)^* \frac{\partial}{\partial (D_t^\alpha \omega)} + \sum_{k=1}^{\infty} (-1)^k D_{d_1} D_{d_2} \dots D_{d_k} \frac{\partial}{\partial (\omega)_{d_1, d_2, \dots, d_k}},$$

where  $(D_t^\alpha)^*$  denotes the adjoint operator for the Riemann-Liouville fractional operator  $(D_t^\alpha)$  and equals to the standard right-hand sided Caputo fractional differential operator [55]. According to [55], the time fractional system of equations (2.5) is said to be nonlinearly self-adjoint whenever by introducing new variables  $f = \psi(x, t, u, v)$  and  $g = \varphi(x, t, u, v)$ , where at least one of them is non-zero and substituting them and their related partial derivatives with respect to  $x$  into adjoint equations (24) and (25), the resultant equations are satisfied for all solutions of the governing problem (1). It means that for nonlinear self-adjointness of (1) the following conditions should be held :

$$\frac{\delta L}{\delta u} = \mu_1 \left( \frac{\partial^\alpha u}{\partial t^\alpha} + v_x + uu_x + qu_{xx} \right) + \mu_2 \left( \frac{\partial^\alpha v}{\partial t^\alpha} + pu_{xxx} + uv_x + vu_x - qv_{xx} \right),$$

$$\frac{\delta L}{\delta v} = \mu_3 \left( \frac{\partial^\alpha u}{\partial t^\alpha} + v_x + uu_x + qu_{xx} \right) + \mu_4 \left( \frac{\partial^\alpha v}{\partial t^\alpha} + pu_{xxx} + uv_x + vu_x - qv_{xx} \right),$$

where  $\mu_i$ , ( $i = 1, 2, 3, 4$ ) are unknown coefficients. The above relations are given as follows:

$$\begin{aligned} & (D_t^\alpha)^* \psi - u(x, t) (\psi_x + \psi_u u_x + \psi_v v_x) - v(x, t) (\varphi_x + \varphi_u u_x + \varphi_v v_x) \\ & + q (\psi_{xx} + 2\psi_{x,u} u_x + 2\psi_{x,v} v_x + \psi_{u,u} u_x^2 + 2\psi_{u,v} u_x v_x + \psi_{u,v} u_{xx} + \psi_{v,v} v_x^2 + \psi_v v_{x,x}) \\ & - p [\varphi_{x,x,x} + 3\varphi_{u,u,v} u_x^2 v_x + 3\varphi_{v,v,x} v_{xx} v_x + 3\varphi_{u,v,v} u_x v_x^2 + 3\varphi_{u,u,u} u_x u_{x,x} + 3\varphi_{u,v,u} u_{x,x} v_x \\ & + 6\varphi_{x,u,v} u_x v_x + 3\varphi_{u,v} u_x v_{x,x} + 3\varphi_{x,x,u} u_x + 3\varphi_{x,x,v} v_x + 3\varphi_{x,u} u_{x,x} + 3\varphi_{x,v} v_{x,x} \\ & + \varphi_u u_{xxx} + \varphi_v v_{xxx} + 3\varphi_{x,u,u} u_x^2 + 3\varphi_{x,v,v} v_x^2 + \varphi_{u,u,u} u_x^3 + \varphi_{v,v,v} v_x^3] \\ & = \mu_1 \left( \frac{\partial^\alpha u}{\partial t^\alpha} + v_x + uu_x + qu_{xx} \right) + \mu_2 \left( \frac{\partial^\alpha v}{\partial t^\alpha} + pu_{xxx} + uv_x + vu_x - qv_{xx} \right), \end{aligned} \quad (26)$$

and

$$(D_t^\alpha)^* \varphi - (\psi_x + \psi_u u_x + \psi_v v_x) - (\varphi_x + \varphi_u u_x + \varphi_v v_x) u - q (\varphi_{xx} + 2\varphi_{x,u} u_x$$

$$\begin{aligned}
& +2\varphi_{x,y}v_x + \varphi_{u,u}u_x^2 + 2\varphi_{u,y}u_xv_x + \varphi_{u}u_{xx} + \varphi_{v,y}v_x^2 + \varphi_{v}v_{x,x} \\
& = \mu_3\left(\frac{\partial^\alpha u}{\partial t^\alpha} + v_x + uu_x + qu_{xx}\right) + \mu_4\left(\frac{\partial^\alpha v}{\partial t^\alpha} + pu_{xxx} + uv_x + vu_x - qv_{xx}\right). \tag{27}
\end{aligned}$$

Now by balancing the coefficients of the different powers of dependent variables  $u$  and  $v$  and their related partial derivatives in both sides of (26) and (27) a systems of partial differential equations should be concluded. Solving the resultant system of differential equations analytically, the following results is obtained:

$$\begin{aligned}
\mu_i &= 0, \quad i = 1, 2, 3, 4, \\
\psi(x, t, u, v) &= f(x, t) = A, \quad \varphi(x, t, u, v) = g(x, t) = B, \tag{28}
\end{aligned}$$

where  $A$  and  $B$  are free constants. This confirms the nonlinear self-adjointness of the problem (1). Corresponding to each vector field  $V_i = \xi_i \partial_x + \tau_i \partial_t + \eta_i \partial_u + \phi_i \partial_v$  there exist Lie characteristic functions which are defined as follows:

$$W_i^u = \eta_i - \xi_i u_x - \tau_i u_t, \quad W_i^v = \phi_i - \xi_i v_x - \tau_i v_t.$$

So Corresponding to the vector fields (12) for the time fractional Whitham-Broer-Kaup problem the following Lie characteristic functions are concluded, respectively:

$$W_1^u = -\alpha u - \alpha x u_x - 2 t u_t, \quad W_1^v = -2 \alpha v - \alpha x v_x - 2 t v_t, \tag{29}$$

and

$$W_2^u = -u_x, \quad W_2^v = -v_x. \tag{30}$$

Now by using the above constructed Lie characteristic functions, conserved vectors corresponding to the governing problem should be calculated. A vector  $C = (C^x, C^t)$  is said to be a conserved vector for the problem (1) if the following conservation equation is held for it,

$$D_t(C^t) + D_x(C^x)_{(1)} = 0. \tag{31}$$

Clearly in the governing system of fractional partial differential equations (1), all of the partial derivatives with respect to the independent variable  $x$  are of integer-orders, so the  $x$  components of the conserved vectors are obtained based on the following classical forms [52]:

$$C_i^x = (W_i^u \frac{\delta L}{\delta u_x} + D_x(W_i^u) \frac{\delta L}{\delta u_{xx}} + D_x^2(W_i^u) \frac{\delta L}{\delta u_{xxx}}) + (W_i^v \frac{\delta L}{\delta v_x} + D_x(W_i^v) \frac{\delta L}{\delta v_{xx}}). \tag{32}$$

Moreover, an extended formula to compute the  $t$ -component of conserved vector for the time fractional system of equations are introduced as follows [55] :

$$C_i^t = \sum_{k=0}^{n-1} (-1)^k [D_t^{\alpha-1-k}(W_i^u) D_t^k(\frac{\delta L}{\delta (D_t^\alpha u)}) + D_t^{\alpha-1-k}(W_i^v) D_t^k(\frac{\delta L}{\delta (D_t^\alpha v)})]$$

$$-(-1)^n [J(W_i^u, \frac{\partial L}{\partial (D_t^\alpha u)}) + J(W_i^v, \frac{\partial L}{\partial (D_t^\alpha v)})], \quad (33)$$

where  $n = [\alpha] + 1$  and  $J(f, g)$  denotes an integral transform which defined as follows:

$$J(f, g) = \frac{1}{\Gamma(n-\alpha)} \int_0^t \int_t^p \frac{f(x, s)g(x, r)}{(r-s)^{\alpha+1-n}} dr ds.$$

Letting  $A = 1$  and  $B = 1$  and by substituting Lie characteristic functions (29) and (30) into relations (32) and (33), the following conserved vectors for the time fractional problem (1) is obtained.

**Case1.**  $0 < \alpha < 1$ , conserved vectors are given as follows:

$$\begin{aligned} C_1^x &= -u(\alpha u + \alpha x u_x + 2t u_t + 2\alpha v + \alpha x v_x + 2t v_t) - v(\alpha u + \alpha x u_x + 2t u_t) \\ &\quad - q(2\alpha u_x + \alpha x u_{xx} + 2t u_{tx} - 3\alpha v_x - \alpha x v_{xx} - 2t v_{tx}) \\ &\quad - p(3\alpha u_{xx} + \alpha u_{xxx} + 2t u_{txx}) - (2\alpha v + \alpha x v_x + 2t v_t), \\ C_1^t &= \alpha(-I_t^{1-\alpha}(u) + 2I_t^{1-\alpha}(v)) - \alpha x(I_t^{1-\alpha}(u_x) + I_t^{1-\alpha}(v_x)) + 2I_t^{1-\alpha}(t v_t) - 2I_t^{1-\alpha}(t u_t), \\ C_2^x &= u(-u_x - v_x) - v u_x - v_x - q(u_{xx} + v_{xx}) - p u_{xxx}, \\ C_2^t &= -I_t^{1-\alpha}(u_x) - I_t^{1-\alpha}(v_x). \end{aligned} \quad (34)$$

**Case2.**  $1 < \alpha < 2$ , conserved vectors are given as follows:

$$\begin{aligned} C_1^x &= -u(\alpha u + \alpha x u_x + 2t u_t + 2\alpha v + \alpha x v_x + 2t v_t) - v(\alpha u + \alpha x u_x + 2t u_t) \\ &\quad - q(2\alpha u_x + \alpha x u_{xx} + 2t u_{tx} - 3\alpha v_x - \alpha x v_{xx} - 2t v_{tx}) \\ &\quad - p(3\alpha u_{xx} + \alpha u_{xxx} + 2t u_{txx}) - (2\alpha v + \alpha x v_x + 2t v_t), \\ C_1^t &= -\alpha(-D_t^{\alpha-1}(u) + 2D_t^{\alpha-1}(v)) - \alpha x(D_t^{\alpha-1}(u_x) + D_t^{\alpha-1}(v_x)) + 2D_t^{\alpha-1}(t v_t) - 2D_t^{\alpha-1}(t u_t), \\ C_2^x &= u(-u_x - v_x) - v u_x - v_x - q(u_{xx} + v_{xx}) - p u_{xxx}, \\ C_2^t &= -D_t^{\alpha-1}(u_x) - D_t^{\alpha-1}(v_x). \end{aligned} \quad (35)$$

#### 4. Conclusions

In this paper, the Lie group analysis approach is used to study time fractional coupled Whitham-Broer-Kaup equations with Riemann-Liouville derivative operators. Based on the Lie symmetries analysis, the governing system of time fractional partial differential equations is similarity reduced to a system of nonlinear fractional ordinary differential equations in Erdelyi-Kober derivative

sense. The new conservation theorem based on the generalization of fractional Noether operators is used to calculate the conserved vectors and conservation laws of the model successfully.

## R E F E R E N C E S

- [1]. *K B Oldham, J Spanier*, The Fractional Calculus : Theory and Application of Differentiation and Integration to Arbitrary Order, Academic Press, New York, NY, USA, 1974.
- [2]. *K. S. Miller and B. Ross*, An Introduction to Fractional Calculus and Fractional Differential Equations (Wiley, New York, 1993)
- [3]. *L Debnath*, Recent applications of Fractional Calculus to science and engineering. Hindawi Publishing Corporation 54. 2003.
- [4]. *A Loverro*, Fractional Calculus: History, Definitions and Applications for the Engineer. Department of Aerospace and Mechanical Engineering University of Notre Dame Notre Dame, U.S.A, 2004
- [5]. *A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo*, Theory and Applications of Fractional Differential Equations, NorthHolland Mathematics Studies Vol. 204 (Elsevier, Amsterdam, 2006).
- [6]. *Uchaikin Vladimir V.* Fractional derivatives for physicists and engineers. Berlin, Springer;2013.
- [7]. *Diethelm, Kai, Neville J. Ford, and Alan D. Freed*. A predictor-corrector approach for the numerical solution of fractional differential equations. *Nonlinear Dynamics* 29.1 (2002) : 3-22.
- [8]. *Zhu, Li, and Yanxin Wang*. Solving fractional partial differential equations by using the second Chebyshev wavelet operational matrix method. *Nonlinear Dynamics* (2017) : 1-11.
- [9]. *Mirzazadeh, M.* Analytical study of solitons to nonlinear time fractional parabolic equations. *Nonlinear Dynamics* 85.4 (2016) : 2569-2576.
- [10]. *Shivanian, Elyas, and Ahmad Jafarabadi*. An improved spectral meshless radial point interpolation for a class of time-dependent fractional integral equations : 2D fractional evolution equation. " *Journal of Computational and Applied Mathematics* 325 (2017) : 1833.
- [11]. *S Abbasbandy*, An approximation solution of a nonlinear equation with Riemann-Liouville's fractional derivatives by He's variational iteration method, *Journal of Computational and Applied Mathematics*. 207(1)(2007) 53-58.
- [12]. *S Kazem, S Abbasbandy, S Kumar*, Fractional-order Legendre functions for solving fractional-order differential equations, *Applied Mathematical Modelling*. 73(7)(2013) 5498-5510.
- [13]. *S Kumar, D Kumar, S Abbasbandy, MM Rashidi*, Analytical solution of fractional Navier-Stokes equation by using modified Laplace decomposition method, *Ain Shams Engineering Journal*. 5(2)(2014) 569-574.
- [14]. *E Shivanian, S Abbasbandy, M S Alhuthali, H H Alsulami*, Local integration of 2-D fractional telegraph equation via moving least squares approximation, *Engineering Analysis with Boundary Elements*. 56 (2015) 98-105.
- [15]. *H Roohani Ghehsareh, S H Bateni, A Zaghian*, A meshfree method based on the radial basis functions for solution of two-dimensional fractional evolution equation, *Engineering Analysis with Boundary Elements*. 61(2015) 52-60
- [16]. *H Roohani Ghehsareh, A Zaghian, SM Zabetzadeh*, The use of local radial point interpolation method for solving two-dimensional linear fractional cable equation. *Neural Computing and Applications* (2017).

- [17] *Saha Ray, S.* A novel method for travelling wave solutions of fractional Whitham-Broer-Kaup, fractional modified Boussinesq and fractional approximate long wave equations in shallow water. *Mathematical Methods in the Applied Sciences* 38.7 (2015) 1352-1368.
- [18] *S.M. El-Sayed, D. Kaya*, Exact and numerical traveling wave solutions of Whitham-Broer-Kaup equations, *Appl. Math. Comput.* 167 (2005) 1339-1349.
- [19] *Xu, Guiqiong, and Zhibin Li*. Exact travelling wave solutions of the Whitham-Broer-Kaup and Broer-Kaup-Kupershmidt equations. *Chaos, Solitons Fractals* 24.2 (2005) 549-556.
- [20] *Rafei, M., and H. Daniali*. Application of the variational iteration method to the Whitham-Broer-Kaup equations. *Computers, Mathematics with Applications* 54.7 (2007) 1079-1085.
- [21] *Ganji D D, Rokni H B, Sfahani M G, Ganji S S*. Approximate traveling wave solutions for coupled Whitham-Broer-Kaup shallow water. *Advances in Engineering Software* 41.7 (2010) 956-961.
- [22] *Jamshidzadeh S, Abazari R*, Solitary wave solutions of three special types of Boussinesq equations, *Nonlinear Dynamics*, 2017.
- [23] *Y. Wang, Yu-Feng Zhang, Zhen-Jiang Liu and Muhammad Iqbal*, A fractional Whitham-Broer-Kaup equation and its possible application to tsunami prevention, *Thermal Science*, 21(4) (2017) 1847-1855.
- [24] *Wang, Linjun, and Xumei Chen*. Approximate analytical solutions of time fractional Whitham-Broer-Kaup equations by a residual power series method. *Entropy* 17.9 (2015) 6519-6533.
- [25] *M D Khan, I Naeem, M Imran*, Analytical solutions of time space fractional, advection-dispersion and Whitham-Broer-Kaup equations, *PRAMANA - Journal of physics*, 83 (6) (2014) 885-906.
- [26] *Amjad Ali, Kamal Shah, Rahmat Ali Khan*, Numerical treatment for traveling wave solutions of fractional Whitham-Broer-Kaup equations, *Alexandria Engineering Journal*, (2017).
- [27] *P.J. Olver*, Application of Lie groups to Differential Equations, Springer-Verlag, (1986).
- [28] *G.W. Bluman, S. Kumei*, Symmetries and Differential Equations, Springer-Verlag, (1989).
- [29] *G.W. Bluman, A.F. Cheviakov, S.C. Anco*, Applications of Symmetry Methods to Partial Differential Equations, Springer, (2010).
- [30] *V. Dorodnitsyn*, Applications of Lie Groups to Difference Equations, CRC Press, (2011).
- [31] *E. Buckwar, Y. Luchko*, Invariance of a partial differential equation of fractional order under the Lie group of scaling transformations, *J. Math. Anal. Appl.* 227 (1998) 81-97.
- [32] *Luchko, Yuri, and Rudolf Gorenflo*. - Scale-invariant solutions of a partial differential equation of fractional order. *Fract. Calc. Appl. Anal.* 3.1 (1998) : 63-78.
- [33] *R.K. Gazizov, A.A. Kasatkin, S.Y. Lukashchuk*, Continuous transformation groups of fractional differential equations, *Vestnik USATU* 9(21) (2007) 125-135.
- [34] *Gazizov, R. K., A. A. Kasatkin, and S. Yu Lukashchuk*. Symmetry properties of fractional diffusion equations. *Physica Scripta* 2009.T136 (2009) : 014016.
- [35] *Hashemi, M. S.* Group analysis and exact solutions of the time fractional Fokker-Planck equation. *Physica A : Statistical Mechanics and its Applications* 417 (2015) : 141-149.
- [36] *Hashemi, M. S., and D. Baleanu*. On the time fractional generalized Fisher equation : group similarities and analytical solutions. *Communications in Theoretical Physics* 65.1 (2016) : 11.
- [37] *Hashemi, M. S., F. Bahrami, and R. Najafi*. Lie symmetry analysis of steady-state fractional reaction-convection-diffusion equation. *Optik-International Journal for Light and Electron Optics* 138 (2017) : 240-249.
- [38] *Hashemi, M. S., and D. Baleanu*. Lie symmetry analysis and exact solutions of the time fractional gas dynamics equation. *JOURNAL OF OPTOELECTRONICS AND ADVANCED MATERIALS* 18.3-4 (2016) : 383-388.

[39] *GW Wang, MS Hashemi*, Lie symmetry analysis and soliton solutions of time-fractional  $K(m,n)$  equation, *Pramana*, 88 (1) (2017).

[40] *R Najafi, F Bahrami, MS Hashemi*, Classical and nonclassical Lie symmetry analysis to a class of nonlinear time-fractional differential equations, *Nonlinear Dynamics*, 87 (3) (2017) 1785-1796.

[41] *S Pashayi, MS Hashemi, S Shahmorad*, Analytical lie group approach for solving fractional integro-differential equations, *Communications in Nonlinear Science and Numerical Simulation*, 51 (2017) 66-77.

[42] *Singla, Komal, R. K. Gupta*. On invariant analysis of some time fractional nonlinear systems of partial differential equations. I. *Journal of Mathematical Physics* 57.10 (2016) 101504.

[43] *Singla, Komal, R. K. Gupta*. Comment on Lie symmetries and group classification of a class of time fractional evolution systems [J. Math. Phys. 56, 123504 (2015)], *Journal of Mathematical Physics* 58, 054101 (2017).

[44] *Singla, Komal, R. K. Gupta*. On invariant analysis of space-time fractional nonlinear systems of partial differential equations. II. *Journal of Mathematical Physics* 58.5 (2017) : 051503.

[45] *Singla, Komal, R. K. Gupta*. Generalized Lie symmetry approach for fractional order systems of differential equations. III, *Journal of Mathematical Physics* 58, 061501 (2017).

[46] *A.V. Bocharov, V.N. Chetverikov, S.V. Duzhin N.G. Chorkova A.M. Vinogradov*, *Symmetries and Conservation Laws for Differential Equations of Mathematical Physics*, 1999.

[47] *E. Noether*, *Invariante Variationsprobleme*, *Knigliche Gesellschaft der Wissenschaften zu Gttingen, Nachrichten. Mathematisch-Physikalische Klasse Heft 2* (1918) 235-257, English transl. : *Transport Theory Statist. Phys.* 1 (3) (1971) 186-207.

[48] *Agrawal OP*. Formulation of Euler-Lagrange equations for fractional variational problems. *Math Anal Appl* 2002; 272 : 368-79.

[49] *Frederico, G.S.F., Torres, D.F.M.* : A formulation of Noethers theorem for fractional problems of the calculus of variations. *J. Math. Anal. Appl.* 334, 834-846 (2007)

[50] *Atanackovic, T.M., Konjik, S., Pilipovic, S., Simic, S.* : Variational problems with fractional derivatives : invariance conditions and Noethers theorem. *Nonlinear Anal.* 71, 1504-1517 (2009)

[51] *Bourdin L, Cresson J, Greff I*. A continuous/discrete fractional Noethers theorem. *Commun Nonlinear Sci Numer Simul* 2013; 18 : 878-887.

[52] *Ibragimov, N.H.* : A new conservation theorem. *J. Math. Anal. Appl.* 333, 311-328 (2007)

[53] *Lukashchuk, S.Y.* : Conservation laws for time-fractional subdiffusion and diffusion-wave equations. *Nonlinear Dyn.* 80, 791-802 (2015)

[54] *Gazizov, R.K., Ibragimov, N.H., Lukashchuk, S.Y.* : Nonlinear self-adjointness, conservation laws and exact solutions of time-fractional Kompaneets equations. *Commun. Nonlinear Sci. Numer. Simul.* 23, 153-163 (2015)

[55] *Singla, Komal, R. K. Gupta*. Conservation laws for certain time fractional nonlinear systems of partial differential equations, *Communications in Nonlinear Science and Numerical Simulation*, 53 (2017) 10-21.

[56] *Singla, Komal, R. K. Gupta*. Space-time fractional nonlinear partial differential equations : symmetry analysis and conservation laws, *Nonlinear Dyn* 2017.

[57] A. Majlesi, H.Roohani Ghehsareh, A. Zaghian, On fractional Jaulent-Miodek equation associated with energy-dependent Schrodinger potential: Lie symmetry reductions, explicit exact solutions and conservation laws, *European journal of physical plus*,

[58] *Kiryakova, V.* : *Generalized Fractional Calculus and Applications*, Pitman Research Notes in Mathematics Series. Longman Scientific Technical, Longman Group, Harlow (1994)